








INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K	A2	(11) International Publication Number: WO 00/04866 (43) International Publication Date: 3 February 2000 (03.02.00)
<p>(21) International Application Number: PCT/US99/16630</p> <p>(22) International Filing Date: 23 July 1999 (23.07.99)</p> <p>(30) Priority Data: 60/093,869 23 July 1998 (23.07.98) US</p> <p>(71) Applicant (for all designated States except US): SOUTHERN RESEARCH INSTITUTE [US/US]; 2000 Ninth Avenue South, Birmingham, AL 35205 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SECRIST, John, A., III [US/US]; 5564 Surrey Lane, Birmingham, AL 35243 (US). TIWARI, Kamal, N. [US/US]; 3045 Brook Highland Drive, Birmingham, AL 35242 (US). MONTGOMERY, John, A. [US/US]; 565 Rutherford Circle, Birmingham, AL 35206 (US).</p> <p>(74) Agents: AMERNICK, Burton, A. et al.; Pollock, Vande Sande & Amernick, Suite 800, 1990 M Street, N.W., Washington, DC 20036 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>

(54) Title: PREPARATION OF THIOARABINOFURANOSYL COMPOUNDS AND USE THEREOF



(57) Abstract

Patients suffering from cancer are treated by being administered a compound represented by formula: (1), wherein each R individually is H or an aliphatic or aromatic acyl group; A is selected from the group consisting of formulae (IIa, IIb, IIc, IIId), wherein X is selected from the group consisting of hydrogen, fluorine, alkoxy, alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, amino, monoalkylamino, dialkylamino, cyano and nitro. The above compounds also inhibit DNA replication in mammalian cells.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PREPARATION OF THIOARABINOFURANOSYL COMPOUNDS AND USE THEREOF

DESCRIPTION

Federally Sponsored Research and Development

5 This invention was supported by Grant CA-34200 from National Institutes of Health.

Technical Field

10 The present invention is concerned with treating patients suffering from cancer by administering to the patients certain thioarabinofuranosyl compounds. Compounds employed according to the present invention have exhibited good anticancer activity. Compounds employed according to the present invention are in the beta configuration as contrasted to the alpha configuration, which failed to exhibit anticancer activity. The present invention also relates to inhibiting DNA replication in a mammalian cell by contacting with the thioarabinofuranosyl compounds. The present invention also relates to a new process for preparing the compounds employed according to the present invention.

Background of Invention


20 Vast amounts of research have accrued over the years related to developing treatments against cancers to inhibit and kill tumor cells. Some of this research has resulted in achieving some success in finding clinically approved treatments. Nevertheless, efforts continue at an ever-increasing rate in view of the extreme difficulty in uncovering promising anticancer treatments. For example, even when

a compound is found to have cytotoxic activity, there is no predictability of it being selective against cancer cells.

One particular compound that has been used rather widespread is cytosine arabinoside, commonly referred to as Ara-C.

Summary of Invention


It has been found according to the present invention that certain thioarabinofuranosyl cytosine compounds are suitable as anticancer agents. The presence of the thio sugar moiety surprisingly makes it possible to achieve good antitumor activity. More particularly, the present invention relates to treating a mammalian host in need of an anticancer treatment by administering to the host an effective anticancer amount of at least one compound represented by the following formula 1:





1


wherein each R individually is H or an aliphatic or aromatic acyl group;

A is selected from the group consisting of





and



5 X is selected from the group consisting of hydrogen, fluoro, alkoxy, alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, amino, monoalkylamino, dialkylamino, cyano and nitro.

10 It has also been found according to the present invention that the above-disclosed compounds of formula 1 can be used to inhibit DNA replication in a mammalian cell by contacting the cell with at least one of these compounds.

The present invention is also concerned with a process for preparing the above-identified compounds. The compounds employed according to the present invention can be prepared by:

A) reacting a 2,3,5-tri-O-aryl or alkyl-4-xylose diaryl or dialkyl dithioacetal such as 2,3,5-tri-O-benzyl-L-xylose-dibenzyl dithioacetal in the presence of a leaving group at the 4 hydroxyl to produce the corresponding 1,4-dithio-D-arabinofuranoside such as benzyl 2,3,5-tri-O-benzyl-1,4-dithio-D-arabinofuranoside;

5 B) subjecting the product from step A) to acidolysis to form the corresponding O-acetyl-4-thio-D arabinofuranose such as 2,3,5-tri-O-benzyl-1-O-acetyl-4-thio-D-arabinofuranose;

10 C) reacting the product of step B with a cytosine, a 5- or 6-aza compound or a suitably blocked derivative thereof forming a corresponding 4-thio- α,β -D-arabinofuranosyl) compound such as, in the case of cytosine, 1-(2,3,5-tri-O-benzyl-4-thio- α,β -D-arabinofuranosyl) cytosine;

D) converting the compound of step C) by hydrolysis to the corresponding thio sugar derivative such as 1-(4-thio- α,β -D-arabinofuranosyl) cytosine;

15 E) separating out the α form of the anomeric mixture of step D) to thereby obtain the desired compound of formula 1, such as 1-(4-thio- β -D-arabinofuranosyl) cytosine.

20 Still other objects and advantages of the present invention will become readily apparent by those skilled in the art from the following detailed description, wherein it is shown and described only the preferred embodiments of the invention, simply by way of illustration of the best mode contemplated of carrying out the invention. As will be realized the invention is capable of other and different
25 embodiments, and its several details are capable of modifications in various obvious respects, without departing from the invention. Accordingly, the description contained herein is to be regarded as illustrative in nature and not as restrictive.


Summary of Drawings

Figs. 1a and 1b are graphs showing metabolism of 2-deoxycytidine, 2'-deoxythiocytidine, ara c and thio ara-c to their respective triphosphates.

Fig. 2 is a graph showing the retention of ara c triphosphate and thio ara c triphosphate in CEM cells.





Best and Various Modes for Carrying Out Invention

The present invention is related to treating a mammalian host in need of an anticancer treatment, which comprises administering to the host an effective anticancer amount of at least compound represented by the formula 1:



Each R in formula 1 individually is preferably H or an aliphatic or aromatic acyl group. Typical aliphatic acyl groups contain from 1 to 6 carbon atoms and include formyl, acetyl, and propionyl. Typical aromatic acyl groups include unsubstituted and alkyl substituted aromatic groups containing 7-10 carbon atoms in the aromatic group. When substituted, the alkyl group typically contains 1-6 carbon atoms. Typical aromatic acyl groups include benzoyl and para-toloyl.

A in formula 1 is preferably



5 Suitable monoalkylamino groups for X contain 1-6 carbon atoms and include monomethylamino, monoethylamino, mono-isopropylamino, mono-n-propylamino, mono-isobutyl-amino, mono-n-butylamino and mono-n-hexylamino. The alkyl moiety can be straight or branched chain.

10 Suitable dialkylamino groups for Y and X contain 1-6 carbon atoms in each alkyl group. The alkyl groups can be the same or different and can be straight or

branched chain. Examples of some suitable groups are dimethylamino, diethylamino, ethylmethylamino, dipropylamino, dibutylamino, dipentylamino, dihexylamino, methylpentylamino, ethylpropylamino and ethylhexylamino.

Suitable halogen groups for X include Cl, Br and F.

5 Suitable alkyl groups for X typically contain 1-6 carbon atoms and can be straight or branched chain. Some examples are methyl, ethyl, i-propyl, n-propyl, i-butyl, n-butyl, pentyl and hexyl.

10 Suitable haloalkyl groups typically contain 1-6 carbon atoms and can be straight or branched chain and include Cl, Br or F substituted alkyl groups including the above specifically disclosed alkyl groups.

 Suitable alkoxy groups typically contain 1-6 carbon atoms and include methoxy, ethoxy, propoxy and butoxy.

 Suitable alkenyl groups typically contain 2-6 carbon atoms and include ethenyl and propenyl.

15 Suitable haloalkenyl groups typically contain 1-6 carbon atoms and include Cl, Br or F substituted alkenyl groups including the above specifically disclosed alkenyl groups.

 Suitable alkynyl groups typically contain 1-6 carbon atoms and include ethynyl and propynyl.

20 The preferred compound employed according to the process of the present invention is 1-(4-thio- β -D-arabinofuranosyl) cytosine.

The present invention is suitable for treating mammalian hosts including humans suffering from cancer including melanoma, prostate cancer, mammary cancer, renal cancer, colon cancer, lung cancer, leukemias and lymphomas.


5 The compounds employed according to the present invention can be prepared by reacting a 2,3,5-tri-O-aryl or alkyl-4-xylose diaryl or dialkyl dithioacetal such as 2,3,5-tri-O-benzyl-L-xylose-dibenzyl dithioacetal in the presence of a bearing group at the 4 hydroxyl position to produce the corresponding 1,4-dithio-D-arabinofuranoside such as benzyl 2,3,5-tri-O-benzyl-1,4-dithio-D-arabinofuranoside. This step can be carried out using phosphine, iodine and imidazole. The product
10 from the above step is subjected to acidolysis to form the corresponding O-acetyl-4-thio-D arabinofuranose such as 2,3,5-tri-O-benzyl-1-O-acetyl-4-thio-D-arabinofuranose. For instance, acetic acid in the presence of mercuric acetate can be employed.

15 The product of the above step is reacted with a cytosine, a 5- or 6-aza compound or a suitably blocked derivative thereof forming a corresponding 4-thio- α,β -D-arabinofuranosyl compound such as, in the case of cytosine, 1-(2,3,5-tri-O-benzyl-4-thio- α,β -D-arabinofuranosyl) cytosine. Suitable blocked derivatives include acyl and trimethylsilylated derivatives. The compound of the above step is converted by hydrolysis to the corresponding thio sugar derivative such as 1-(4-thio- α,β -D-arabinofuranosyl) cytosine.
20

The α form of the anomeric mixture of the above step is separated out to thereby obtain the desired compound of formula 1, such as 1-(4-thio- β -D-arabinofuranosyl) cytosine.

25 Compounds according to the present invention can be prepared by the process sequence shown in scheme 1 and examples 1 and 2 described hereinbelow, wherein the preferred compound, 1-(4-thio- β -D-arabinofuranosyl) cytosine is illustrated for purposes of facilitating an understanding of the process sequence. The precursor employed, 2,3,5-thio-O-benzyl-L-xylose dibenzyl dithioacetal, can be produced by the process described by Secrist, III et al, "The Synthesis and

Biological Activity of Certain 4'-Thionucleosides, *Nucleosides & Nucleotides*, 14 (3-5), 675-686 (1995), disclosure of which is incorporated herein by reference. The process of the present invention provides for a relatively efficacious method for producing significant quantities of the desired compound. Prior art techniques are extremely complicated and not readily suitable for producing desired amounts of the compounds.



The pharmaceutically acceptable effective dosage of the active compound of the present invention to be administered is dependent on the species of the warm-blooded animal (mammal), the body weight, age and individual condition, and on the form of administration.

5 The pharmaceutical composition may be oral, parenteral, suppository or other form which delivers the compounds used in the present invention into the bloodstream of a mammal to be treated.

10 The compounds of the present invention can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but generally administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

15 The dosage administered will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the age, health and weight of the recipient; the nature and extent of the symptoms, the kind of concurrent treatment; the frequency of treatment; and the effect desired. A daily dosage of active ingredient can be expected to be about 0.001 to 1000 milligram (mg) per kilogram (kg) of body weight, with the preferred dose being 0.1 to about 30 mg/kg.

20

Dosage forms (compositions suitable for administration) typically contain from about 1 mg to about 100 mg of active ingredient per unit. In these pharmaceutical compositions, the active ingredient will ordinarily be present in an amount of about 0.5-95% by weight based on the total weight of the composition.

5 The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, and powders, or in liquid dosage forms, such as elixirs, syrups, and suspensions. It can also be administered parenterally, in sterile liquid dosage forms. The active ingredient can also be administered intranasally (nose drops) or by inhalation. Other dosage forms are potentially possible such as administration transdermally, via a patch mechanism or ointment.

10 Gelatin capsules contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar-coated or film-coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

15 Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

20 In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water-soluble salt of the active ingredient, suitable stabilizing agents, and, if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propylparaben, and chlorobutanol.

25 Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences. Mack Publishing Company, a standard reference text in this field.

Useful pharmaceutical dosage forms for administration of the compounds according to the present invention can be illustrated as follows:

Capsules

5 A large number of unit capsules are prepared by filling standard two-piece hard gelatin capsules each with 100 mg of powdered active ingredient, 150 mg of lactose, 50 mg of cellulose, and 6 mg of magnesium stearate.

Soft Gelatin Capsules

10 A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil, or olive oil is prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 mu of the active ingredient. The capsules are washed and dried.

Tablets

15 A large number of tablets are prepared by conventional procedures so that the dosage unit was 100 mg of active ingredient, 0.2 mg of colloidal silicon dioxide, 5 mg of magnesium stearate, 275 mg of microcrystalline cellulose, 11 mg of starch, and 98.8 mg of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

20 Various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

The foregoing disclosure includes all the information deemed essential to enable those skilled in the art to practice the claimed invention. Because the cited applications may provide further useful information, these cited materials are hereby incorporated by reference in their entirety.

5

The following non-limiting examples are presented to further illustrate the present invention.

Example 1**Preparation of 1-(4-Thio- β -D-Arabinofuranosyl) Cytosine**

2,3,5-Tri-O-benzyl-L-xylose Dibenzy l Dithioacetal (4). L-Xylose (1, 25g, 167 mmol) was stirred for 5 hours in 0.5% hydrogen chloride in methanol (675 mL) at room temperature and then neutralized with Amberlite IRA-400 OH anion exchange resin. The filtrate and washings were combined and evaporated to dryness and the crude product was purified by silica gel chromatography ($\text{CHCl}_3/\text{MeOH}$, 92:8) to afford 26.2 g of methyl L-xylofuranoside (2, 95% yield) as an α and β (1:1) mixture. MS 164 (M)⁻, 165 ($\text{M}+\text{H}$)⁺, 133 ($\text{M}-\text{OCH}_3$)⁺.

To an ice-cold solution of 2 (10 g, 60.9 mmol) in dry tetrahydrofuran (350 mL) was added sodium hydride (60% dispersion in mineral oil, 14.8 g, 370 mmol) and the reaction mixture was stirred for 15 min under N_2 . To this reaction mixture was added solid tetrabutylammonium iodide (0.36 g, 0.96 mmol) followed by a dropwise addition of benzyl bromide (36.6 g, 214 mmol). The reaction mixture was stirred for 3 days at room temperature. After the addition of methanol (25 mL) the solution was evaporated under reduced pressure, and the crude product was purified by silica gel chromatography (cyclohexane/EtOAc, 9:1) to afford pure methyl 2,3,5-tri-O-benzyl-L-xylofuranoside (3, 23 g, 87% yield). MS 435 ($\text{M}+\text{H}$)⁺, 433 ($\text{M}-\text{H}$)⁺, 403 ($\text{M}-\text{OCH}_3$)⁺; ^1H NMR (CDCl_3) δ 7.38-7.25 (m, 30H, aromatic $\text{H}=\text{s}$), 4.94 (d, 1H, $\text{H}-1\alpha$, $J_{1,2} = 4.3$ Hz), 4.87 (d, 1H, $\text{H}-1\beta$, $J_{1,2} = 0.9$ Hz), 4.64-4.45 (m, 12H, PhCH_2 's), 4.37 (m, 1H, $\text{H}-4\alpha$), 4.27 (dt, 1H, $\text{H}-4\beta$, $J_{4,5a} = 3.7$ Hz, $J_{4,5b} = 6.5$ Hz, $J_{3,4} = 6.2$ Hz), 4.17 (t, 1H, $\text{H}-3\alpha$, $J_{3,4} = 6.9$ Hz, $J_{2,3} = 5.6$ Hz), 4.07 (dd, 1H, $\text{H}-3\beta$, $J_{3,4}$

= 6.2 Hz, $J_{2,3} = 2.5$ Hz), 4.00 (dd, 1H, H-2 α , $J_{2,3} = 5.6$ Hz), 3.95 (t, 1H, H-2 β , $J_{2,3} = 2.5$ Hz), 3.70 (dd, 1H, H-5a α , $J_{4,5a} = 4.5$ Hz, $J_{5a,5b} = 10.4$ Hz), 3.66 (dd, 1H, H-5a β , $J_{4,5a} = 3.7$ Hz, $J_{5a,5b} = 10.7$ Hz), 3.54 (dd, 1H, H-5b α , $J_{4,5b} = 7.5$ Hz), 3.49 (dd, 1H, H-5b β , $J_{4,5b} = 6.5$ Hz).

5 To a solution of **3** (42 g, 97 mmol) in dichloromethane (1000 mL) were added benzyl mercaptan (49.6 g, 400 mmol) and stannic chloride (4.93 g, 18.9 mmol), and the reaction mixture was stirred at room temperature overnight. After neutralization with 5% aqueous NaHCO₃ (750mL), the organic layer was separated and the aqueous layer was extracted with dichloromethane (500 mL). The combined organic layers were
10 evaporated, and crude **4** was purified by silica gel chromatography (cyclohexane/EtOAc, 99:1) to afford **4** (8.53 g, 57%) of sufficient purity to carry forward. MS 657 (M+Li)⁺; ¹H NMR (CDCl₃) δ 7.35-7.29 (m, 19H, aromatic *H*=s), 7.19-7.13 (m, 4H, aromatic *H*=s), 7.01-6.96 (m, 2H aromatic *H*=s), 4.86 (d, 1H, PhCHH, $J = 11.1$ Hz), 4.70 (two overlapping d=s, 2H, PhCHH, PhCHH, $J = 11.1$ Hz, $J = 11.2$ Hz), 4.43 (d, 1H, PhCHH, 11.2 Hz), 4.40 (d, 1H, PhCHH, $J = 11.9$ Hz), 4.36 (d, 1H, PhCHH, $J = 11.9$ Hz), 4.07 (dd, 1H, H-2, $J_{1,2} = 3.0$ Hz, $J_{2,3} = 7.5$ Hz), 3.75-3.67 (m, 4H, two PhCH₂=s), 3.68 (d, 1H, H-1, $J_{1,2} = 3.0$ Hz), 3.36-3.25 (m, 2H, H-4, H-5a), 3.15-3.12 (m, 1H, H-5b), 2.22 (d, 1H, 4-OH, $J = 6.2$ Hz).

2,3,5-Tri-*O*-benzyl-1-*O*-acetyl-4-thio-D-arabinofuranose (6). To a solution of 4 (13.0 g., 20 mmol) in dry 2:1 toluene/acetonitrile (200 mL) were added triphenylphosphine (15.7 g, 60 mmol), iodine (12.7 g, 50 mmol) and imidazole (5.44 g, 80 mmol). The reaction mixture was stirred at 90 °C for 24 h after which time the solution was evaporated to dryness. The crude product was purified by silica gel chromatography (cyclohexane/EtOAc, 4:1) to afford benzyl 2,3,5-tri-*O*-benzyl-1,4-dithio-D-arabinofuranoside as a syrup (5, 9.0 g, 83%). MS 543 (M+H)⁺; ¹H NMR (CDCl₃) δ 7.40-7.20 (m, 20H, aromatic *H*=s), 4.69-4.42 (m, 6H, three PhCH₂O=s), 4.37 (m, 1H, H-1), 4.20 (m, 2H, H-2, H-3), 3.87 (s, 2H, PhCH₂S-), 3.80 (dd, 1H, H-5a, J_{4,5a} = 7.4 Hz, J_{5a,5b} = 9.3 Hz), 3.55 (dd, 1H, H-5b, J_{4,5b} = 7.1 Hz), 3.47 (m, 1H, H-4). Anal. (C₃₃H₃₄O₃S₂ @ 0.25 H₂O) C, H.

To a suspension of mercuric acetate (7.29 g, 22.9 mmol) in acetic acid (96 g) was added 5 (5.42 g, 10 mmol), and the resulting mixture was stirred at room temperature for 2 h. The reaction mixture was diluted with dichloromethane (200 mL) and washed successively with water, saturated aqueous NaHCO₃ and 5% aqueous KCN solution. The organic layer was dried over Na₂SO₄ and concentrated. Chromatography of the crude product using cyclohexane:ethylacetate (98:2) as eluent gave a mixture of α and β (1:1) anomers of 6 (3.73 g, 78%) as a colorless syrup. MS 479 (M+H)⁺; ¹H NMR (CDCl₃) δ 7.35-7.23 (m, 15H, aromatic *H*=s), 6.07 (d, 0.25H, H-1β, J_{1,2} = 4.0 Hz), 5.98 (d, 0.75H, H-1α, J_{1,2} = 2.8 Hz), 4.83-4.45 (m, 6H, PhCH₂'s), 4.26 (dd, 0.75H, H-2α, J_{2,3} = 5.4 Hz), 4.17-4.11 (m, 0.5H, H-2β, H-3β), 4.03 (t, 0.75H, H-3α, J_{3,4} = 6 Hz), 3.80-3.67 (m, 1.25H, H-4α, H-5aα, H-5aβ), 3.53-3.39 (m, 1.75H, H-5bα, H-4β, H-5bβ), 2.06 (s, 3H, CH₃-α and CH₃-β). Anal. (C₂₈H₃₀O₅S @ 0.75 H₂O) C, H.

1-(2,3,5-Tri-*O*-benzyl-4-thio- α,β -D-arabinofuranosyl) cytosine (7). To a suspension of 1-*O*-acetyl 2,3,5-tri-*O*-benzyl-4-thio-D-arabinofuranose (478 mg, 1 mmol) and cytosine (111.0 mg, 1 mmol) in anhydrous acetonitrile (25 mmol) were added consecutively hexamethyldisilazane (HMDS, 162 mg, 1 mmol) and chlorotrimethylsilane (TMSCl, 434 mg, 4 mmol), and the mixture was stirred at room temperature for 0.5 hours. This solution was cooled to -78 °C. Trimethylsilyltrifluoromethane sulfonate (267 mg, 1.2 mmol) was added and the resulting solution was stirred at -78 °C for another 2.5 h, after which time the reaction was essentially complete. The mixture was warmed to room temperature, concentrated to a small volume (5 mL), diluted with methylene chloride (50 mL) and then washed with water (20 mL) followed by saturated sodium bicarbonate and water. The organic layer was dried over MgSO₄ and evaporated to dryness. The residue was purified by chromatography over silica gel (50 g. elution with CHCl₃/MEOH 98:2) to afford 7 (412 mg 77.5%) as a colorless syrup TLC (95:5 CHCl₃/MEOH) R_f 0.65; MS z/e 536 (M+Li)⁺.

¹H NMR (CDCl₃) δ 8.22 (d, 1, H-6 _{β} , J = 7.6); (d, 1, H-6 _{α} , J_{5,6} = 7.5 Hz); 7.38-7.09 (m, 30, aromatic H's); 6.65 (d, 1, H-1' _{β} , J_{1',2'} = 5.7 Hz); 6.36 (d, 1, H-1' _{α} , J_{1',2'} = 1.2 Hz); 5.44 (d, 1, H-5 _{α}); 5.26 (d, 1, H-5 _{β}); 4.97-4.33 (overlapping multiplets, 12, C₆H₅CH₂); 4.26 (dd, 1 H-2' _{β} , J_{2',3'} = 6.8 Hz); 4.22 (m, 1, H-3' _{α} , J_{3',4'} = 1 Hz); 4.16 (dd, 1, H-3' _{β} , J_{3',4'} = 6.4 Hz); 4.13 (m, 1, H-2' _{α} , J_{2',3'} = 1.7 Hz); 3.91 (m, 1, H-4' _{α}); 3.78 (m, 1, H-5' _{α}); 3.73-3.63 (m, 2, H-5' _{β}); 3.55 (m, 1, H-5' _{α}); 3.46 (m, 1, H-4' _{β}).

1-(4-Thio- α , β -D-arabinofuranosyl) cytosine (8). To a solution of boron trichloride (1 M solution) in dry dichloromethane (7mL, 7mmol) cooled to -78 °C, was added dropwise over a period of 30 min, a solution of compound 7 (265 mg, 0.5 mmol) in dry dichloromethane (10 mL). Stirring was continued overnight at -20 °C. The solvent was removed in vacuo and the residue was coevaporated with dichloromethane (4 x 20 mL). The residue was neutralized with saturated NaHCO₃ (25 mL) and washed with chloroform (15 mL). The aqueous layer was applied to a cation exchange (H⁺) column and eluted with water to remove salts and then eluted with 1N NH₄OH to obtain the desired compound 8 (110 mg, 85%) MS z/e 260(M+H)⁺.

¹H NMR (Me₂SO-*d*₆) δ 7.94 (d, .67, H-6- β , J_{5,6} = 7.5 Hz); 7.90 (d, 0.33, H-6 α , J_{5,6} = 7.5 Hz); 7.17-7.03 (overlapping bs, 2, NH's); 6.33 (d, 0.67, H-1' β , J_{1',2'} = 4.6 Hz); 5.86 (d, 0.33, H-1' α , J_{1',2'} = 7.3 Hz); 5.77 (d, 0.33, H-5 α); 5.70 (d, 0.67, H-5 β); 5.61 and 5.59 (overlapping doublets, 1, 2'-OH β , J_{2',2'-OH} = 5.1 Hz, 2'-OH α , J_{2',2'-OH} = 5.9 Hz); 5.47 (d, 0.33, 3'-OH α , J_{3',3'-OH} = 5.1 Hz); 5.38 (d, 0.67, 3'-OH β , J_{3',3'-OH} = 4.2 Hz); 5.08 (t, 0.67, 5'-OH β , J_{5',5'-OH} = 5.4 Hz); 4.90 (t, 0.33, 5'-OH α , J_{5',5'-OH} = 5.2 Hz); 4.00-3.93 (m, 1.67, H-2' α , H-2' β , H-3' β); 3.86-3.76 (m, 1, H-5' α and H-5' β); 3.67-3.55 (m, 1, H-5' β + H-3' α); 3.49-3.33 (m, 0.67, H-4' α , H-5' ω); 3.17 (m, 0.67, H-4' β).

1-(5-*O*-dimethoxytrityl-4-thio- β -D-arabinofuranosyl) cytosine (9). To a solution of compound 8 (100 mg, 0.38 mmol) in dry pyridine (10 mL) was added 4,4'-Dimethoxytritylchloride (135 mg, 0.6 mmol) and the solution was stirred for 2 h at room temperature. The reaction mixture was evaporated to dryness and the crude was dissolved in ethyl acetate (20 mL) and washed with water and evaporated to dryness to afford a solid which was purified on silica gel column (CHCl₃/MeOH 98:2) to obtain pure compound 9 (96 mg, 90% based on 1:1 mixture of α , β -mixture of 8). MS *m/z* 568.3 (M+Li)⁺.

¹H NMR (Me₂SO-*d*₆) δ 7.77 (d, 1, H-6, *J*_{5,6} = 7.5 Hz); 7.42-7.23 (m, 9, aromatic H's); 7.17 (bs, 1, NH); 7.05 (bs, 1, NH); 6.91-6.88 (m, 4, aromatic H's); 6.36 (d, 1, H-1', *J*_{1',2'} = 4.8 Hz); 5.65 (d, 1, H-5); 5.76 (d, 1, 2'-OH, *J*_{2',2'-OH} = 4.6 Hz); 5.43 (d, 1, 3'-OH, *J*_{3',3'-OH} = 3.3 Hz); 3.98-3.91 (m, 2, H-2', H-3'); 3.75 (s, 6, OCH₃); 3.39-3.25 (m, 4, H-4', H-5').

1-(4-Thio- β -D-arabinofuranosyl) cytosine (10). Compound 9 (90 mg, 0.16 mmol) was treated with trifluoroacetic acid (22 mg) in chloroform (5 mL) at room temperature for 10 min. The reaction mixture was neutralized with aq. NaHCO₃ and the aqueous layer was applied on a cation exchange column which was eluted first with water to remove salt and finally with 1N NH₄OH to afford compound 10 (35 mg, 85%), mp 218-220 °C (lit¹ 221-222 °C); MS z/e 260(M+H)⁺.

¹H NMR (Me₂SO-*d*₆) δ 7.94 (d, 1, H-6, J_{5,6} = 7.5 Hz); 7.12 (bs, 1, NH); 7.04 (bs, 1, NH); 6.33 (d, 1, H-1', J_{1',2'} = 4.6 Hz); 5.70 (d, 1, H-5); 5.61 (bd, 1, 2'-OH, J_{2',2'-OH} = 3.1 Hz); 5.38 (bd, 1, 3'-OH, J_{3',3'-OH} = 3.5 Hz); 5.08 (bt, 1, 5'-OH, J_{5',5'-OH} = 4.9 Hz); 4.00-3.93 (m, 2, H-2', H-3'); 3.78 (m, 1, H-5'_a); 3.61 (m, 1, H-5'_b); 3.16 (m, 1, H-4').

Example 2

Preparation of 1-(4-Thio- β -D-Arabinofuranosyl)5-Fluorocytosine

1-(2,3,5-Tri-*O*-benzyl-4-thio- α,β -D-arabinofuranosyl)5-fluorocytosine (7a). To a suspension of 1-*O*-acetyl 2,3,5-tri-*O*-benzyl-4-thio-D-arabinofuranose (478 mg, 1 mmol) and cytosine (129.0 mg, 1 mmol) in anhydrous acetonitrile (25 mmol) were added consecutively hexamethyldisilazane (HMDS, 162 mg, 1 mmol) and chlorotrimethylsilane (TMSCl, 434 mg, 4 mmol), and the mixture was stirred at room temperature for 0.5 hours. This solution was cooled to -78°C. Trimethylsilyltrifluoromethane sulfonate (267 mg, 1.2 mmol) was added and the resulting solution was stirred at -78°C for another 2.5 h, after which time the reaction was essentially complete. The mixture was warmed to room temperature, concentrated to a small volume (5 mL), diluted with

methylen chloride (50 mL) and then washed with water (20 mL) followed by saturated sodium bicarbonate and water. The organic layer was dried over MgSO_4 and evaporated to dryness. The residue was purified by chromatography over silica gel (50 g. elution with $\text{CHCl}_3/\text{MEOH}$ 98:2) to afford **7** as 2:1 α,β mixture (425.2 mg 80.0%) as a colorless syrup TLC (95:5 $\text{CHCl}_3/\text{MEOH}$ R_f 0.65; MS z/e 553 ($\text{M}+\text{Li}$)⁺.

¹H NMR (CDCl_3) δ 8.40 (d, 1, H-6_B, $J=7.6$); 8.10 (d, 1, H-6_A, $J_{5,6} = 7.5$ Hz); 7.38-7.09 (m, 30, aromatic H's); 6.55 (d, 1, H-1'_B, $J_{1',2'} = 5.7$ Hz); 6.25 (d, 1, H-1'_A, $J_{1',2'} = 1.2$ Hz); 4.90 (d, 1, H-5_A); 4.78 (d, 1, H-5_B); 4.30-4.55 (overlapping multiplets, 12, $\text{C}_6\text{H}_5\text{CH}_2$); 4.15 (dd, 1 H-2'_B, $J_{2',3'} = 6.8$ Hz); 4.10 (m, 1, H-3'_A, $J_{3',4'} = 1$ Hz); 3.90 (dd, 1, H-3'_B, $J_{3',4'} = 6.4$ Hz); 3.75 (m, 1, H-2'_A, $J_{2',3'} = 1.7$ Hz); 3.70 (m, 1, H-4'_A); 3.65 (m, 1, H-5'_A); 3.50-3.55 (m, 2, H-5'_B); 3.45 (m, 1, H-5'_B); 3.40 (m, 1, H-4'_B).

1-(4-Thio- β -D-arabinofuranosyl)5-fluorocytosine (10a). To a solution of boron trichloride (1 M solution) in dry dichloromethane (7 mL, 7 mmol) cooled to -78°C , was added dropwise over a period of 30 min, a solution of compound **7** (2273 mg, 0.5 mmol) in dry dichloromethane (10 mL). Stirring was continued overnight at -20°C . the solvent was removed in vacuo and the residue was coevaporated with dichloromethane (4 x 20 mL). The residue was neutralized with saturated NaHCO_3 (25 nL) and washed with chloroform (15 mL). The aqueous layer was applied to a cation exchange (H^+) column and eluted with water to remove salts and then eluted with 1N NH_4OH to obtain the desired compound **8a** as 2:1 α,β mixture which upon crystallization with water gave pure **10a** (32.2 mg, 25%) MS z/e 278($\text{M}+\text{H}$)⁺,

¹H NMR ($\text{Me}_2\text{SO}-d_6$) δ 8.15 (d, 1, H-6, $J_{5,6} = 7.5$ Hz); 7.75 (bs, 1, NH); 7.50 (bs, 1, NH); 6.25 (d, 1,

H-1', $J_{1',2'} = 4.6$ Hz); 5.65 (d, 1, H-5); 5.60 (bd, 1, 2'-OH, $J_{2',2'-OH} = 3.1$ Hz); 5.40 (bd, 1, 3'-OH, $J_{3',3'-OH} = 3.5$ Hz); 5.20 (bt, 1, 5'-OH, $J_{5',5'-OH} = 4.9$ Hz); 4.00-3.90 (m, 2, H-2', H-3'); 3.75 (m, 1, H-5'_a); 3.65 (m, 1, H-5'_b); 3.15 (m, 1, H-4').

5 ¹ Ototani, H.; Whistler, R.L. Preparation and Antitumour Activity of 4'-Thio Analogs of 2,2'-Anhydro-1-β-D-arabinofuranosylcytosine. *J. Med. Chem.* **1974**, *17*, 535-537.

Example 3

The following in vitro tests were carried out.

Table 1
IC₅₀ Molar Values

	ThioaraC	5-fluorothioaraC	Cytosine arabinoside
LOX IMV1 Melanoma	7×10^{-6}	6×10^{-6}	1×10^{-6}
PC-3 Prostate	8×10^{-6}	1×10^{-4}	5×10^{-6}
ZR-75-1 Mammary	3×10^{-6}	6×10^{-7}	4×10^{-7}
CAK1-1 Renal	3×10^{-6}	2×10^{-5}	6×10^{-7}
DLD-1 Colon	7×10^{-6}	8×10^{-5}	5×10^{-6}
NC1-H23 Lung	5×10^{-6}	3×10^{-6}	3×10^{-7}
SK-MEL 28 Melanoma	7×10^{-6}	—	2×10^{-6}
SNB-7 (CNS)	1×10^{-6}	4 x	2×10^{-7}

The above tests were carried out with a 72 hour exposure to the compound using Natural Red assay for all cell lines except SK-MEL 28 which was done with SRB assay.

Example 4

The following *in vivo* tests were carried out to illustrate the effectiveness of the compounds of the present invention.

Table 2
Response of SC Colon 36 Tumor to Treatment

Treatment				Tumor Regression						
Agent	Dose Mg/Kg	Rt	Schedule	Non-Specific Deaths/ Total	Number of Partial	Number of Complete	Duration Med/Range (Days)	Tumor Free Surv./Total	Days to 2 Doubling	Days Delay (T-C)
Control		IP	Q 4HX 3 Day 9-17					0/10	10.5	
1-(4-thio-β-D-arabinofuranosyl)cytosine (prepare in saline/ injection volume .5 cc/20g. body weight	8	IP	Q 4HX 3 day 9-17	0/5	0	4		4/5	>49.0	>38.5
Palmo-ARA-C (prepared in saline/injection volume .5cc/20 g body weight	18	IP	Q 1DX 9 day 9	0/5	0	5	34.6 35-35	4/5	>49.0	>38.5

Palmo-ARA-C is a prodrug of ARA-C.

Table 3
Response of SC CAKI-1 Renal Tumor to Treatment

Treatment				Tumor Regression						
Agent	Dose mg/kg	Rt	Schedule	Non-Specific Deaths/ Total	Number of Partial	Number of Complete	Duration Med/Range (Days)	Tumor Free Surv/Total	Days to 3 Doubling	Days Delay (T-C)
Control		IP	Q 4 HX 3 day 7-15					0/12	9.5	
1-(4-thio-8-D-arabino-furanosyl) cytosine	16	IP	Q 4HX 3 day 7-15	1/6	0	5	26.9 16-48	0/6	59.8	50.3
1-(4-thio-8-D-arabino-furanosyl) cytosine prepared in saline; injection volume: 0.5cc/20 g body weight	8	IP	Q 4 HX 3 day 7-15	0/6	1	4	12.7 9-18	0/6	50.7	41.2
ARA-C	30	IP	Q 4HX 3 day 7-15	5/6	0	0		0/6	25.7	16.2 (Toxic)
ARA-C	20	IP	Q 4HX 3 day 7-15	5/6	0	0		0/6	15.7	6.2 (Toxic)
ARA-C prepared in saline; injection volume: 0.1cc/10g body weight	13.3	IP	Q 4HX 3 day 7-15	0/6	0	0		0/6	20.5	11.0
Palmo-ARA-C	45	IP	Q 1DX 5 day 7	4/6	0	0		0/6	41.2	31.7 (Toxic)
Palmo ARA-C prepared in saline/tween 80; injection volume 0.5cc/20g body weight	22.5	IP	Q 1DX 9 day 7	1/6	0	1	9.1 9-9	0/6	36.0	26.5

Table 4
Response of SC HCT-116 Colon Tumor to Treatment

Treatment				Tumor Regression						
Agent	Dose Mg/Kg	Rt	Schedule	Non- Specific Deaths/ Total	Number of Partial	Number of Complete	Duration Med/Range (Days)	Tumor Free Surv/Total	Days to 2 Doubling	Days Delay (T-C)
Control		IP	Q 4HX 3 day 6-14					0/12	8.9	
1-(4-thio-8-O-arabino-furanosyl) cytosine	20	IP	Q 4HX 3 day 6-14	0/6	0	6		6/6	>52.0	>43.1
1-(4-thio-8-O-arabino-furanosyl) cytosine prepared in saline (soluble); injection volume 0.5cc/20g body weight	13.3	IP	Q 4HX 3 day 6-14	0/6	0	6	23.6	5/6	>52.0	>43.1
ARA-C	13.3	IP	Q 4HX 3 day 6-14	0/6	0	0	24-24			
ARA-C prepared in saline; injection volume; 0.5cc/20g body weight	9	IP	Q 4HX 3 day 6-14	0/6	0	0		0/6	12.2	3.3
Palmo-ARA-C	22.5	IP	Q 1DX 9 day 6	0/6	0	1	24.2	0/6	24.0	15.1
Palmo-ARA-C prepared in saline + tween 80; injection volume; 0.5cc/20g body weight	15	IP	Q 1DX 9 day 6	0/6	0	0	24-24	0/6	14.9	6.0

* Palmo-ARA-C is a prodrug of ARA-C.

As can be appreciated from the above tests, in most cell lines, ARA-C exhibits higher cytotoxicity as compared to that of the present invention which would tend to discourage use of the compounds of the present invention as a cancer treatment. However, surprisingly, the 1-(4-thio- β -D-arabinofuranosyl)cytosine is more potent than ARA-C and more selective in killing cancer cells *in vivo*. Compounds employed according to the present invention exhibit better anticancer activity along with lower toxicity. This may be due to the fact that 1-(4-thio- β -D-arabinofuranosyl)cytosine does not get deaminated by cytidine deaminase as quickly as AraC does, which is shown below in Table 5 by using radio labeled substances. Good substrate activity of cytidine deaminase with AraC has been a major problem with this drug because it not only decreases half-life of drug but also gets converted to AraUridine which is not an active substance.

Table 5

Cytidine deaminase activity
determined with radioactive substrates

Compound	K_m	V_{max}	V_{max}/K_m
AraC	147	9	0.06
1-(4-thio- β -D-arabinofuranosyl cytosine	2571	17	0.007

The following tests were carried out which illustrate the ability of the thioarabinofuranosyl compounds according to the present invention to inhibit DNA replication in mammalian cells. In the following tests, dCyd refers to 2'-deoxycytidine, TCyd refers to thiodeoxycytidine, T-araC refers to 1-(4-thio- β -D-arabinofuranosyl) cytosine and Ara-C refers to "cytosine arabinoside". The terms "d CTP", "T-d CTP", "ara CTP" and "T ara CTP" refer to the corresponding triphosphates. The term "dURD" refers to 2-deoxyuridine. The term "F-dUrd" refers to fluorinated deoxyuridine and "F-dUMP" refers to fluorinated deoxyuridine monophosphate.

Reference is made to Figs. 1a and 1b which illustrate the metabolism of dCyd, T-dCyd, araC and T-araC to their respective triphosphates. In particular, CEM cells were incubated with 2000 nM of [5-³H]T-dCyd, 5 nM [5-³H]araC, or 25 nM [5-³H]T-araC for the times indicated. The radioactivity in the 5'-triphosphate peak was determined using SAX HPLC. For each compound, the amount of radioactivity in the 5'-triphosphate peak was more than 90% of the total radioactivity eluting from the SAX HPLC column.

Test results related to the inhibition of CEM cell growth are shown below in Table 6.

CEM cells were incubated with various concentrations of each compound and the effect on cell numbers was determined using a Coulter Counter. The IC₅₀ was determined from a plot of the percent of control growth *versus* drug concentration. The data presented are the mean and standard deviation from 3 separate experiments. The araC-resistant cells were obtained by incubating CEM cells in the presence of 150 nM araC for approximately 1 month at which time the CEM cells were growing at control rates.

Table 6
Inhibition of CEM Cell Growth by dCyd Analogs

Compound	Wild-type	araC-resistant	Resistant/WT
IC ₅₀ (nM)			
T-dCyd	2200 ± 1400	7200 ± 7600	3
T-araC	24 ± 9	125 ± 84	5
araC	6 ± 3	800 ± 400	145

Test results related to metabolism in wild-type and araC resistant CEM cells are shown below in Table 7.

CEM cells (wild-type or araC-resistant cells) were incubated with 100 nM of [5-³H]dCyd, [5-³H]T-dCyd, [5-³H]araC, or [5-³H]T-araC for the times indicated.

The radioactivity in the 5'-triphosphate peak and the incorporation of radioactivity in the acid-insoluble fraction (DNA) was determined. For each compound, the amount of radioactivity in the 5'-triphosphate peak was more than 90% of the total radioactivity eluting from the SAX HPLC column.

Table 7
Metabolism of dCyd and Its Analogs in
Wild-Type and araC Resistant CEM Cells

Compound	Incubation time Hours	Cell type	Triphosphate pmoles/10 ⁶ cells (percent of WT)	DNA	Total
dCyd	0.25	WT	2.27	1.19	3.46
		Resistant	0.021 (0.93)	0.011 (0.89)	0.032 (0.91)
T-dCyd	0.25	WT	0.877	1.29	2.17
		Resistant	0.036 (4.1)	0.051 (4.0)	0.087 (4.0)
araC	1	WT	14.7	0.28	14.98
		Resistant	0.19 (1.2)	0.018 (6.3)	0.208 (1.4)
T-araC	1	WT	0.202	0.0056	0.2076
		Resistant	0.051 (25)	0.0025 (45)	0.0535 (26)

Results illustrating phosphorylation in cell-free CEM extracts are shown below in Table 8.

A crude cell extract was prepared from wild-type and araC-resistant CEM cells and the ability to phosphorylate dCyd, araC, and T-araC was determined. The number in parentheses is the number of experiments performed. Reactions were performed in solutions containing 50 mM Tris (pH 8.0), 5 mM ATP, 7.5 mM MgCl₂, 20 mM NaF, extract, and 1 μ M [5-³H]dCyd, [5-³H]araC or [5-³H]T-araC. At various times after the initiation of the experiment, an aliquot of each reaction volume was removed and placed on DE-81 filters. The filters were washed with 1 mM ammonium formate and ethanol, and the amount of radioactivity on each filter was determined. The phosphorylation of each compound increased in a linear manner with respect to time. This assay was verified by HPLC.

Table 8

Phosphorylation of dCyd, araC, and T-araC

in Cell-Free CEM Extracts

Compound	Wild Type	araC-resistant CEM cells	Percent of wild-type
	pmoles/mg protein/minute		
dCyd	240 \pm 33 (3)	3.5 \pm 0.2 (3)	1.5
araC	94 (1)	1.2 (1)	1.3
T-araC	0.83 (2)	0.011 (1)	1.3

Table 9 below illustrates deoxycytidine deaminase activity. In particular, deoxycytidine deaminase activity was purified from Molt-4 cells as described (Shewach et al, *Mol. Pharmacol.* 42: 518-524, 1992). Each assay was done in duplicate and the kinetic constants were determined from linear double-reciprocal plots of 1/velocity versus 1/concentration of the substrate. The best line was

determined by linear regression from at least 5 datum points, and the K_m and V_{max} were determined from the x and y intercepts.

Table 9

Substrate Characteristics of dCyd, araC, and T-araC

with dCyd Kinase Activity Isolated from Molt-4 Cells

Compound	K_m (μM)	Relative V_{max}	V_{max}/K_m
dCyd	1.2	1	0.8
araC	15	0.1	0.006
4'-thio-araC	93	0.46	0.005

Table 10 below also illustrates deoxycytidine deaminase activity.

Deoxycytidine deaminase activity was purified from human placenta and the K_m and V_{max} of dCyd, T-dCyd, araC, and T-araC were determined. Reactions were carried out in solutions containing 20 mM potassium phosphate (pH 7.4), 100 mM KCl, various concentrations of radiolabeled nucleoside, and enzyme. The reactions were stopped with acid, the substrate was separated from the product by paper chromatography, and the radioactivity in each were determined. The kinetic constants were determined from linear double-reciprocal plots of 1/velocity versus 1/concentration of the substrate. The best line was determined by linear regression from at least 5 datum points, and the K_m and V_{max} were determined from the x and y intercepts. The data presented are the mean and standard deviation from 3 separate experiments.

Table 10

Substrate Characteristics of dCyd, T-dCyd, araC, and T-araC

with dCyd Deaminase Activity Isolated from Human Placenta

Compound	K_m (μ M)	V_{max} (pmoles/mg/min)	V_{max}/K_m
dCyd	23 ± 2.7	13 ± 1.5	0.55
T-dCyd	111 ± 77	37 ± 12	0.33
araC	238 ± 131	14 ± 7	0.058
T-araC	2944 ± 1023	21 ± 6	0.0072

Results of half-life tests in CEM cells are reported below in Table 11.

After incubation of CEM cells for 1 hour with 100 nM of [5-³H]dCyd, [5-³H]T-dCyd, [5-³H]araC, or [5-³H]T-araC, the cells were collected, washed with fresh medium, and resuspended in fresh medium that did not contain radiolabeled nucleosides. Samples were collected at various times after the cells were resuspended in fresh medium, and the amount of radioactivity in the 5'-triphosphate peak was determined using SAX HPLC. The data presented are the mean and standard deviation from 3 separate experiments.

Table 11

Initial Half-Life of dCTP, T-dCTP, araCTP,

and T-araCTP in CEM Cells

Nucleotide	Hours (SD)
dCTP	0.94 ± 0.16
T-dCTP	1.10 ± 0.35
araCTP	5.31 ± 0.31
T-araCTP	10.8 ± 1.80

Results of tests related to retention of araCTP and T-araCTP in CEM cells are shown in Fig. 2. The tests were carried out as follows:

After incubation of CEM cells for 1 hour with either 5 nM [5-³H]araC or 200 nM [5-³H]T-araC, the cells were collected, washed with fresh medium, and resuspended in fresh medium that did not contain radiolabeled nucleosides. Samples were collected at various times after the cells were resuspended in fresh medium, and the amount of radioactivity in the 5'-triphosphate peak was determined using SAX HPLC. In this experiment there was 0.639 pmoles araCTP/10⁶ cells and 0.246 pmoles of T-araCTP after the 1 hour incubation with radiolabeled compound.

Results related to metabolism in CEM cells are reported below in Table 12. In particular, CEM cells were incubated with 100 nM of [5-³H]dCyd, [5-³H]T-dCyd, [5-³H]araC, or [5-³H]T-araC for the times indicated. The complete metabolism of each compound was determined: The medium was analyzed for the original compound, its deaminated form using reverse phase HPLC, and H₂O; the acid-soluble extract was analyzed by SAX HPLC for phosphorylated metabolites; and the incorporation of radioactivity into acid-precipitable material (DNA) was determined. All of the original radioactivity was accounted for in these fractions. For each compound, the amount of radioactivity in the 5'-triphosphate peak was more than 90% of the total radioactivity eluting from the SAX HPLC column.

Table 12

Metabolism of dCyd, T-dCyd, araC, and T-araC in CEM Cells

Nucleoside	Time of Incubation	Original Compound	Deaminated Compound	H ₂ O	nucleoside-TP	DNA
	(hours)			(percent of total)		
dCyd	1	38	0	55	4	3
T-dCyd	1	90	0	0	2	8
araC	1	87	0	0	13	0.4
T-araC	24	96	0	0	4	0.3

Results from tests related to the effect of F-dUrd on metabolism are shown below in Table 13.

CEM cells were incubated with 100 nM of [5-³H]dCyd, [5-³H]T-dCyd, [5-³H]araC, or [5-³H]T-araC for the times indicated in the presence or absence of 10 μM F-dUrd, which is metabolized to F-dUMP (a potent inhibitor of thymidylate synthetase). The complete metabolism of each compound was determined as described in the legend to Table 12, and the percent of deaminated metabolites for each compound was determined. In the absence of F-dUrd, the deaminated product for dCyd was [³H]H₂O due to the removal of [³H] at the 5 position of F-dUMP by thymidylate synthetase. In the presence of F-dUrd, the deaminated product of dCyd, T-dCyd, and araC was in the medium as dUrd, T-dUrd, or araU, respectively.

Table 13

Effect of F-dUrd on the Metabolism of

dCyd, T-dCyd, araC, or T-araC

	Compound	F-DUrd (μ M)	Incubation Time (hours)	Percent Deamination
5	dCyd	0	1	32
	dCyd	10	1	22
	T-dCyd	0	8	0
	T-dCyd	10	8	24
	araC	0	8	0
10	araC	10	8	10
	T-araC	0	24	0
	T-araC	10	24	0

Table 14 below reports results of tests concerning effects of DNA, RNA and protein synthesis.

15 CEM cells were incubated with 60 μ M T-dCyd, 150 nM araC, or 750 nM T-araC. Radiolabeled precursors of DNA [3 H]dThd, RNA [3 H]Urd, or protein [3 H]leucine) were added to each treatment 30 minutes after the addition of each compound. Samples were taken 1, 2, 3 and 4 hours after the addition of radiolabeled precursors, and the incorporation of radioactivity into RNA, DNA, or protein were determined as described (Parker et al, *Biochem. Pharmacol.* 55: 1673-1681, 20 1998). Each number represents the average of two experiments.

Table 14

Effect of T-dCyd, araC, and T-araC on

DNA, RNA, and Protein Syntheses

	Compound	DNA	RNA	protein
		Percent of control		
5	60 μ M T-dCyd	38	95	78
	150 nM araC	8	96	96
	750 nM T-araC	16	87	72

As can be appreciated from the above, thioarabinofuranosyl compounds according to the present invention can be used to inhibit DNA replication in mammalian cells. These results tend to suggest their ability as immunomodulators, which would render them suitable for treating autoimmune diseases. This is further supported by the fact that several guanosine analogs have been shown to stimulate the immune system (Weigle, W.O., *CRD Crit.Rev.Immunol.*, 1987, 7, 285; Lin et al., *J.Med.Chem.*, 1985, 28, 1194-1198; Reitz et al., *J.Med.Chem.*, 1994, 37, 3561-3578; Michael et al., *J.Med.Chem.*, 1993, 36, 3431-3436). Certain 3- β -ribofuranosylthiazolo[4,5-d]pyrimidines have also been shown to have significant immunoactivity, including murine spleen cell proliferation and in vivo activity against Semliki Forest virus (Nagahara et al., *J.Med.Chem.*, 1990, 33, 407-415).

The foregoing description of the invention illustrates and describes the present invention. Additionally, the disclosure shows and describes only the preferred embodiments of the invention but, as mentioned above, it is to be understood that the invention is capable of use in various other combinations, modifications, and environments and is capable of changes or modifications within the scope of the inventive concept as expressed herein, commensurate with the above teachings and/or the skill or knowledge of the relevant art. The embodiments described hereinabove are further intended to explain best modes known of

practicing the invention and to enable others skilled in the art to utilize the invention in such, or other, embodiments and with the various modifications required by the particular applications or uses of the invention. Accordingly, the description is not intended to limit the invention to the form disclosed herein. Also, it is intended that the appended claims be construed to include alternative embodiments.

5

CLAIMS



What is claimed is:

1. A method for treating a mammalian host in need of an anticancer treatment which comprises administering to said host an effective anticancer amount of at least one compound represented by the formula 1:




5 wherein each R individually is H, an aliphatic acyl group or an aromatic acyl
6 group;


7 A is selected from the group consisting of



10



11



12

13

14

wherein X is selected from the group consisting of hydrogen, fluorine, alkoxy, alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, amino, monoalkylamino, dialkylamino, cyano and nitro.


1

2. The method of claim 1 wherein each R is H.

1

3. The method of claim 1 wherein A is

2



1

4. The method of claim 3 wherein X is H.

1

5. The method of claim 1 wherein X is H.

6. The method of claim 1 wherein said compound is 1-(4-thio-β-D-arabinofuranosyl) cytosine.


7. The method of claim 1 wherein said host is suffering from a cancer selected from the group consisting of melanoma, prostate cancer, mammary cancer, renal cancer, colon cancer, lung cancer, leukemias and lymphomas.




8. A method for inhibiting replication of DNA which comprises contacting said cell with an effective amount of at least one compound represented by the formula 1:



wherein each R individually is H, an aliphatic acyl group or an aromatic acyl group;

A is selected from the group consisting of






12 wherein X is selected from the group consisting of hydrogen, fluorine,
 13 alkoxy, alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, amino, monoalkylamino,
 14 dialkylamino, cyano and nitro.

1 9. The method of claim 8 wherein each R is H.

1 10. The method of claim 8 wherein A is



11. The method of claim 10 wherein X is H.

1 12. The method of claim 8 wherein X is H.

13. The method of claim 8 wherein said compound is 1-(4-thio- β -D-arabinofuranosyl) cytosine.

1 14. A method for producing a compound of the formula 1 which
2 comprises:

3 A) reacting a 2,3,5-tri-O-aryl or alkyl-4-xylose diaryl or dialkyl
4 dithioacetal in the presence of a leaving group at the 4 hydroxyl position to produce
5 the corresponding 1, 4-dithio-D-arabinofuranoside;

6 B) reacting said arabinofuranoside from step A) to acidolysis to form the
7 corresponding O-acetyl-4-thio-D arabinofuranose;

8 C) reacting said O-acetyl-4-thio-D-arabinofuranose from step B) with a
9 compound selected from the group consisting of a cytosine, a 5-aza compound, a 6-
10 aza compound and blocked derivatives thereof to form the corresponding 4-thio- α,β -
11 D-arabinofuranosyl compound;

12 D) converting said arabinofuranosyl compound from step C) by
13 hydrolysis to the corresponding thio sugar derivative; and

14 E) separating out the α form of the anomeric mixture of step D) to
15 thereby obtain the compound of formula 1.

1 15. The method of claim 14 wherein said compound in step C) is
2 cytosine.

1 16. The method of claim 14 wherein said hydrolysis in step D) comprises
2 reacting said arabinofuranosyl compound from step C) with boron trichloride.

1 17. The method of claim 14 wherein step E) comprises reacting the
2 racemic mixture from step D) with 4,4'-dimethoxytritylchloride to thereby produce
3 5'-O-dimethoxytrityl-4'-thio- α - β -D-arabinofuranosylcytosine, separating 5'-O-
4 dimethoxytrityl-4'-thio- β -D arabinofuranosylcytosine; and then reacting 5'-O-
5 dimethoxytrityl-4'-thio- β -D arabinofuranosylcytosine with trifluoroacetic acid to
6 thereby obtain the compound of formula 1.

1 18. The method of claim 14 wherein said dithioacetal compound in step
2 A) comprises 2,3,5-trio-O-benzyl-xylose dibenzyl dithioacetal.

1 19. The method of claim 14 wherein said dithioacetal is obtained by
2 reacting L-xylose with methanol to produce methyl L-xylofuranoside and reacting
3 said methyl L-xylofuranoside with benzyl bromide.

1/2




FIG. 1A




FIG. 1B

2/2




FIG. 2